



---

## Influence of Telemedicine and Its Dimensions on Health Promotion

<sup>1\*</sup>Mahycol Reynner Harold Bravo-Ramirez, <sup>2\*</sup>Danis Yanet Castillo-Esquivel, <sup>3\*</sup>Josefa Magdalena Cabrera Díaz

<sup>1</sup>E-mail: mbravor11@ucvvirtual.edu.pe

Lecturer of the Faculty of Medical Sciences, Universidad Cesar Vallejo, Trujillo, Peru

<https://orcid.org/0000-0003-4038-0776>

<sup>2</sup>E-mail: danis.castillo@upn.pe

Lecturer of the Faculty of Humanities, Universidad Privada del Norte, Lima, Peru

<https://orcid.org/0000-0001-8811-3154>

<sup>3</sup>Lecturer of the Faculty of Health Sciences, Universidad Privada Antenor Orrego, Trujillo, Peru

E-mail: jcabrerad@upao.edu.pe

<https://orcid.org/0000-0002-4593-7046>

\*These authors contributed equally to this work

Correspondent author:

Mahycol Reynner Harold Bravo-Ramírez

Calle Mantaro 517, Trujillo, Perú

Telephone: +51 949495747

E-mail: mbravor11@ucvvirtual.edu.pe

*(Received: 02 September 2023*

*Revised: 14 October*

*Accepted: 07 November)*

---

### KEYWORDS

Health Promotion,  
Telemedicine,  
Teleconsultation,  
Teleorientation,  
Telemonitoring.

### ABSTRACT:

**Introduction:** Multidisciplinary health promotion activities were interrupted during the pandemic with telemedicine temporarily replacing face-to-face patient care as an alternative. **Objective:** The objective of the research was to determine the influence of Telemedicine and its dimensions on health promotion. **Method:** The type of study is cross-sectional. It was carried out in a public hospital where 120 users participated during the period September - November 2022, using instruments validated by experts and having as inclusion criteria the patients attended in the four remote services: Teleconsultation, Teleorientation, Telemonitoring and Teleinterconsultation, being analyzed statistically by correlation coefficient and logistic regression. **Results:** A response rate of 76.4% was obtained, in which 58.3% presented a high level of satisfaction with the use of Telemedicine, showing a positive influence on health promotion ( $r=0.669$ ,  $p<0.05$ ; OR 9.28 [95 % CI, 4.54 - 18.98;  $p<0.05$ ]). **Conclusion:** It was determined that there is a relationship between Telemedicine and its dimensions towards health promotion as well as high satisfaction on the part of the users.

## I. INTRODUCTION

Public health is in permanent evolution according to the situation of each country, focusing on aspects related to health promotion, prevention and interventions. In this sense, it involves the care of the individual as a unique human being and as part of a society. In addition, in the 21st century, healthcare systems coexist with various aspects such as population aging, migration to cities, global competitiveness,

demand for information management, population beliefs, and economic and/or budgetary constraints[1]. In this sense, public health has been undermined at the beginning of 2020 as a consequence of the global crisis that all countries have faced due to the SARS-CoV-2 virus infection, since the health crisis that has arisen has made visible the limitations of the different health systems, which was evidenced in the collective health component and in the health care services that highlighted the inequities in the social and economic



determinants, especially in Latin American countries[2]. In addition, human resource gaps, such as shortages of medical supplies, biomedical equipment and medicines, were also revealed [3].

In this sense, the different governments had the need to implement and/or reinforce strategies that would allow health personnel to continue providing care to users of the different health services. To this end, they adopted various remote care modalities, making use of Information and Communication Technologies (ICT) and several other services, including Telemedicine, which is a modality designed to ensure that health care can continue to be provided to users, thus avoiding the need for them to visit health facilities in person and thereby prevent the spread of the virus [4]. This is an unexpected challenge for public health, since the health policies of the different governments have been faced with unknown events and existing resources, which have revealed shortcomings in the health structures at both the legal and regulatory levels [3], for example, in the USA, face-to-face care decreased due to the use of Telemedicine [5]. For all these reasons, the COVID 19 pandemic has made public health -oriented in its purposes aimed at promoting health and preventing disease- transcend beyond its statutes and concepts, incorporating into this new scenario regulations and statutes that regulate the actions of health professionals [6].

In Latin America, one of the main supports to the health system, which was immersed in this health situation, is Telehealth, which although it is true that in several countries it was regulated with some guidelines in the areas of application, it took off due to the mentioned pandemic and is being used until today. Its implementation was due to the need to continue providing health services in contrast to the different health measures that were implemented to curb the incidence of new cases, such as social distancing and home isolation. Thus, various modalities of remote health care were implemented by publishing various legislative and regulatory decrees, with the aim of guaranteeing that the population would continue to receive health care considering their health diagnoses and/or avoiding complications,[1,7] Within this framework, the Law No. 30421 was passed in Peru, which establishes general provisions related to the National Telehealth Network, detailing the development axes that

define Telemedicine, which is considered to be the provision of health-oriented services.[8]

In Peru, the modality of Telemedicine care covers 4 dimensions according to the Regulation of Law No. 30421, Framework Law of Telehealth [8], where Teleconsultation is defined as the remote service in which health personnel attends a user to provide a diagnosis within the framework of their skills and competences. This requires the use of ICTs by the users to optimally achieve the objectives pursued, such as health promotion, disease prevention, diagnosis, treatment, rehabilitation and palliative care, as appropriate. Likewise, it is also related to the issue of drug treatment in accordance with the limits of prescription of drugs according to the competence of health professionals or others provided by the Ministry of Health. Whereas, Teleorientation encompasses the set of actions where health professionals use ICTs to provide health counseling and advice to health users with the aim of promoting health, preventing disease, supporting recovery and reducing sequelae. In the same sense, Telemonitoring encompasses the permanent monitoring and/or follow-up by health professionals of users who suffer from chronic diseases and, therefore, are already aware of their diagnosis and treatment. For this purpose, clinical information and medical history is of great importance as it is based on medical standards, biomedical parameters and/or ancillary tests as a means of monitoring their health. Likewise, medication prescriptions may or may not be included depending on medical judgment and the competence of other healthcare professionals. Finally, Teleinterconsultation is the remote consultation requested by a health professional to another specialist, for the attention of a specific health case, in which users may or may not be present. This will make it possible to reconsider the medication treatment or also the performance of auxiliary tests that allow an optimal treatment according to the norms stipulated by the Ministry of Health.

In this sense, Telemedicine is emerging as a sustainable health strategy, proposed by the United Nations in the framework of the 2019-2030 Agenda [9]. Its purpose is to remove the access barrier for users living in distant locations to health facilities, avoid unnecessary travel, strengthen the use of human and logistical resources, allowing their efficient use. In addition, it allows the population to receive care from home, reduces



health gaps and obstacles related to health inequalities, making it possible to guarantee a healthy life and the well-being of the population [1], in such a way that coordination, continuity and integration of the different health systems can be improved, as well as coordinated work between general practitioners and specialists. Moreover, as in other health professionals and the participation of patients in primary care [10], it will allow better control of chronic and metabolic diseases, which in users who suffer from them represent a high risk of mortality due to COVID-19 [11].

In Peru, being a developing country, together with its geographical distribution, health systems are mostly made up of first level primary care facilities and one of their main functions is health promotion, which is a fundamental part of the strengthening of the health systems. Another purpose is to respond to the health needs of individuals, families and communities considering a high level of solidarity and equity, covering not only clinical services from primary care, but also assessing the living conditions and social environment by establishing connections with health professionals [9]. Therefore, knowing that health promotion is an essential function of public health and should be integrated into all aspects related to health processes, this research decided to determine the level of satisfaction, as well as the influence of telemedicine, considering its four dimensions towards the promotion of health in patients treated in public hospitals in the city of Trujillo.

## II. METHODS

### Description Of The Design

The design consists of a cross-sectional, retrospective study conducted in a hospital in northern Peru, in which activities are carried out at different levels of prevention. The data collected covered the period from September to November 2022, which was obtained through telephone calls and the completion of virtual surveys through online digital platforms, as well as access to digitized medical records. This will provide demographic data, user care services, diagnosis codes and other characteristics that will allow us to determine whether the inclusion criteria have been met.

### Population

The study included 120 health users out of a total of 157 (response rate 76.4%), to which a non-randomized probability sample was applied, grouped according to the outpatient services in which they were attended, and considering as inclusion criteria patients over 18 years of age, of both sexes, who had been provided with the four remote care services covered by Telemedicine: Teleconsultation, Teleguidance, Telemonitoring and Teleinterconsultation. On the other hand, participants with cognitive and/or mental deficits and who had not been treated under the 4 telemedicine modalities were excluded.

### Instruments

In relation to the application of the questionnaires, the approval of the hospital's ethics committee was obtained; likewise, telephone calls were made to users and informed consent was requested by means of virtual forms sent through digital tools. For the validation of the questionnaires, a pilot sample of 30 people (not included in the study) was used for both variables, obtaining an overall Cronbach's alpha of 0.89, in addition to being validated by 5 telemedicine experts. The questionnaires for both variables were formulated using a Likert scale with the following values: never = 1, almost never = 2, sometimes = 3, almost always = 4, always = 5, with high scores ranging from 74 to 100 points, regular 47 to 73 points and low scores for those below 47 points. Regarding the telemedicine dimensions, each one was represented by 5 items each, with a Likert scale mentioned above, being stratified from 5 to 11 points as low, 12 to 18 as regular and 19 to 25 as high.

### Statistical Analysis

Participants were stratified by age group and gender, described by a frequency table and percentages, while age was shown as mean and standard deviation.

The normality of the distribution was performed using the Kolmogorov-Smirnov test. The numerical variables according to their distribution that were used were Pearson's correlation test (age - telemedicine) and Spearman's correlation test (health promotion and telemedicine) with  $p < 0.05$ , being considered statistically significant.



The logistic regression model was estimated to evaluate the association between sociodemographic variables and the dimensions of Telemedicine with health promotion. All statistical tests were performed at a 5% significance level with a 95% confidence interval. Microsoft Office Excel was used for data entry and SPSS V.26.0 was used for statistical analysis. The research was conducted in accordance with the ethical principles of the Declaration of Helsinki.

### III. RESULTS

The univariate analysis, as shown in Table 1, demonstrated greater participation of the female gender in 69.2%; Teleconsultation and teleorientation dimensions had high satisfaction representing 77.5% and 65.8% respectively. On the other hand, the dimension that obtained the highest percentage in relation to low satisfaction was the Teleinterconsultation dimension with 17.5%, represented repeatedly with 81% by women. At the Telemedicine level, a average score of 74.79 (SD  $\pm$ 17.97) was obtained, with a high level of satisfaction in 58.3%, represented mostly by women (71.4%) and those under 55 years of age (67.5%), whereas 33.3% and 83% rated it as fair and low, respectively.

Regarding the health promotion variable, half of the participants considered it to be good, with 60% of the female sex being represented; whereas, the rest of the participants rated it as regular and low satisfaction with 42.5% and 7.5%, respectively.

TABLE I: Sociodemographic characteristics of patients and satisfaction levels of Telemedicine dimensions.

Variable	Categories	Number of Cases	%	Mean (SD)
Age (years)	< 55	83	69,2	45,37 (15,27)
	55-64	26	21,7	
	>64	11	9,2	
Gender	Female	83	69,2	
	Male	37	30,8	
Teleconsultation	Low	4	3,3	

	Regular	23	19,2	20.93
	High	93	77,5	(4.44)
Teleorientation	Low	9	7,5	19.77 (5)
	Regular	32	26,7	
	Alto	79	65,8	
Telemonitoring	Low	18	15	17.88 (5.4)
	Regular	48	40	
	High	54	45	
Teleinterconsultation	Low	21	17,5	16.42 (4.9)
	Regular	54	47	
	High	45	37,5	

SD: Standard Deviation

A Pearson correlation was applied between age and Telemedicine, finding an inverse but not significant relationship ( $r$ : -0.076,  $p$ >0.05). Similarly, a Spearman correlation was performed between health promotion and Telemedicine ( $r$ =0.669,  $p$ <0.05), being a strong and significant correlation. In the same way, it was evidenced between the dimensions of Teleconsultation ( $r$ =0.639,  $p$ <0.05), Teleorientation ( $r$ =0.634,  $p$ <0.05), Telemonitoring ( $r$ =0.587,  $p$ <0.05) and Teleinterconsultation ( $r$ =0.605,  $p$ <0.05). In the characteristics of patients associated with level of Telemedicine satisfaction in multivariate analysis showed that compared to age, older than 64 years, those of younger age were less associated with the degree of satisfaction in Telemedicine (younger than 55 years: OR 0.68 [95 % CI, 0.16 -2.9;  $p$ >0.05]; 55-64 years: OR 0.77 [95 % CI, 0.1 - 3.7;  $p$ >0.05]). Women were significantly associated with the degree of satisfaction with Telemedicine use: (OR 1.27 [95 % CI, 0.59 -2.73;  $p$ <0.05]).

Likewise, the association between the variables of our study, such as Telemedicine and health promotion (dependent variable) was determined by logistic regression, which showed a significant association between the two: OR 9.28 [95 % CI, 4.54 - 18.98;  $p$ <0.05].



TABLE II: Association of sociodemographic variables and telemedicine dimensions with the level of health promotion satisfaction through logistic regression.

Variable	Level	OR	Confidence interval at 95%		Value P
			Inferior	Superior	
Gender (ref = Male)	Female	0.460	0.208	1.018	0.055
	Age (ref = over 64 years old)	under 55 years old	0.606	0.165	2.220
	55-64 years	0.537	0.128	2.257	0.396
Telemedicine (Dimensions)	Teleconsultation	15.834	5.163	48.560	0.000 *
	Teleorientation	11.437	5.150	25.399	0.000 *
	Telemonitoring	5.769	3.098	10.741	0.000 *
	Teleinterconsultation	5.371	2.887	9.992	0.000 *

\*:p<0.05

On the other hand, sociodemographic variables (sex, age) and dimensions of Telemedicine were related to the level of satisfaction with health promotion, as shown in Table 2, which is indicative that male gender is related to a higher degree of satisfaction with health promotion, OR 0.46 [95 % CI, 0.20 - 1.01; p<0.05] despite not being significant. Likewise, ages younger than 64 years were not associated with the level of satisfaction with health promotion (younger than 55 years: OR 0.60 [95 % CI, 0.16 - 2.22; p>0.05]; 55-64 years: OR 0.53 [95 % CI, 0.12 - 2.25; ; p>0.05]), not being significant. On the contrary, significant associations were shown with each of the dimensions, highlighting Teleconsultation OR 15.83 [95 % CI, 5.16 - 48.56; p<0.05].

#### IV. DISCUSSION

There is little knowledge about the application of Telemedicine in the different stages of health promotion [12], since to date no studies have been found where the relationship between both variables is mentioned. Currently, international studies focus Telemedicine on activities that are separate from health care and health promotion. Similarly, few studies were found indicating the repercussions of Telemedicine and its dimensions towards health promotion, despite the fact that some authors mention that 50% of Telemedicine use occurs in primary care activities [13].

On the other hand, the findings show that more than half of the patients have a high level of satisfaction, similar to other studies [14–16]. On the other hand, there are studies in which face-to-face care generates greater satisfaction due to the nature of the disease, in which the user necessarily requires direct attention from the health professional [17], which is supported by a North American study that showed that there is no difference between face-to-face care and telemedicine [18].

An influential factor in telemedicine care was gender, for example in Brazil, where it was shown that men are the biggest users of telemedicine [19], in contrast to our country, where women predominate (69.2%), as corroborated by several studies [20–22]. This difference occurs in our research because it encompasses all health promotion activities, while in Scofano et al. [19], it was in home care for patients undergoing hemodialysis. Another influential factor was age, which was in agreement with young adults, and which was associated with greater use of telemedicine, with a representativeness of around one third of the participants [20,22,13].

The association between satisfaction with the Teleconsultation and its influence on satisfaction with health promotion was also determined ( $r=0.639$ ,  $p<0.05$ ; OR 15.834 [95 % CI, 5.163 - 48.560;  $p<0.05$ ]). This is verified with other studies focused on prevention in which prenatal visits were made by telephone, generating satisfaction in almost all the participants [23]. Likewise, if it focuses on other specialties applied to Telemedicine, such as Teledermatology, where it is compared with face-to-face care, indicating the similarity, thus generating high satisfaction between the two [24]. In



addition, studies have been carried out, which include two healthcare centers comparing the degree of satisfaction, thus generating a satisfaction rate of over 60% in both [25], having even mentioned the privacy generated by remote care compared to face-to-face care, being an important factor for its acceptance, considering additionally a lower cost and efficiency in time, especially for adults [26].

On the other hand, there are systematic reviews on the effectiveness of Teleconsultation in primary care where it is considered an effective alternative; however, it brings with it a high attrition rate and cannot be applicable to all [27]. This result is corroborated by the study carried out in Peru, a country with ethnic diversity and in the process of development, where there is a high percentage of rural areas with limited access to Internet and health services, generating a negative social impact since the use of ICTs is limited to the use of text messaging and direct telephone calls, without making use of videoconferencing applications. In addition, there is the slow progress in the implementation of broadband communication networks, generating negative effects on the quality of health care [28]. Therefore, some studies have shown that there is no significant effect at the time of intervening in lifestyles, as in the case of nutritional control both in primary (home) care and in the follow-up of discharged patients, since they consider that they present a risk similar to that of patients who do not receive care [29,30].

Regarding Teleorientation, given in primary care as in the case of Galán's study [31], carried out in a medical post where their main activity is prevention and health promotion, they showed that one fifth of the participants presented low satisfaction, while 36.6% considered it high, being different from our result (65.8%), being surpassed by other international studies where they presented up to 95.3% satisfaction as in oncology patients [32], as well as activities of prevention and control of blood pressure in pregnant women [33]. A limitation in relation to Teleorientation is that there are not many studies that evaluate this dimension; moreover, in all the selected studies the sample was homogeneous and the sample size was small.

Telemonitoring, as is known, is applicable to diagnosis or clinical follow-up through laboratory tests and treatment. This is supported by studies which conclude that it is an accessible

tool especially in primary care for various age groups, especially when directed at older adults and in the control and treatment of patients with chronic diseases, such as hypertension and diabetes mellitus, finding improvements in glucose levels between 6 to 12 months of control [34,35], even mentioning that there was no difference between face-to-face care and telemonitoring in diabetic patients [36], in addition to the participation of multidisciplinary personnel such as pharmaceutical chemists who contribute to initiate and adjust medications according to the patient's needs in order to improve the quality of care, in addition to telemanagement, which is addressed by chemists as they are implementing telemedicine, which is already a reality in several countries [37].

On the other hand, although it is considered an accessible tool, its effectiveness decreases in some specific contexts, such as the deficiency of Telepharmacy in urban areas, especially in the delivery of medicines to oncology patients [16]. More difficulty is encountered in rural areas, due to the series of barriers presented by professionals, since when applying telemonitoring, factors such as technological literacy, knowledge of the use of technology by patients, as well as age and beliefs, must be considered [34,38,39]. Even in hospital activity, this dimension does not have a positive impact on patients infected by COVID 19, since although they were previously monitored by pulse oximetry, they did not have a positive influence on hospital readmissions, hospital stay and the mortality generated by this pandemic. On the contrary, they presented a 12% increase in emergency room attendance and 24% increase in admission to the intensive care unit [40,41].

Teleinterconsultation is a dimension that has been little addressed by international research, and a Peruvian study showed 47.7% satisfaction with this dimension, although it has a negative evaluation in different aspects such as the "structure" dimension, which shows a lack of training and acceptance on the part of health professionals. In addition, other barriers are connectivity as well as aspects of the process itself, where the time of attention is insufficient, since the maximum time of attention is considered to be 8 minutes (a standard time of attention in Peru is between 15 to 20 minutes per patient). Finally, on the outcome aspect, as mentioned above, this is limited to text messages either by vis SMS or



WhatsApp, which is associated with connectivity limitations [42]. A similar problem is experienced in Brazil where, in addition to the aforementioned problems, multiculturalism intervenes in the doctor/patient relationship as they consider the implementation of an informed consent for the exposure of the patient to more than one health professional. For this task, it is difficult to gather more than one at the same time for the care of a patient due to the professionals' schedules [43].

## V. LIMITATIONS

We recognize that this study has limitations, particularly because the study involves data collected from a single hospital. This restricts generalization due to the sample size, and the inclusion criteria that the users had to meet, since they had to be attended by means of the 4 dimensions of Telemedicine, even though no study has been found in which these dimensions of health promotion are addressed in a single investigation, thus limiting absolute comparisons. This study also fails to address individual-level barriers for the use of Telemedicine in rural populations.

## VI. CONCLUSIONS

It is concluded that Telemedicine and its dimensions are related and have a positive influence on health promotion, especially in the satisfaction generated by its use. Likewise, it is necessary to eliminate the multiple barriers that exist for health users, from the political to the technological, in order to be able to cover care throughout the country without any restriction.

## FINANCIAL SUPPORT

This research is self-financed by the authors.

## CONFLICT OF INTEREST

There is no conflict of interest.

## BIBLIOGRAPHY

- [1] Saigí-Rubió, F.; Torrent-Sellens, J.; Robles, N.; Pérez Palaci, J. E.; Baena, M. I. Estudio Sobre Telemedicina Internacional En América Latina: Motivaciones, Usos, Resultados, Estrategias y Políticas. *Estud. sobre telemedicina Int. en América Lat. Motiv. usos, Result. estrategias y políticas* **2021**. <https://doi.org/10.18235/0003438>.
- [2] Suarez, V.; Cabezas, C. Amenazas a La Salud Pública En El Perú y La Respuesta Del Instituto Nacional de Salud. *Rev. Peru. Med. Exp. Salud Publica* **2022**, *39* (2), 140–142. <https://doi.org/10.17843/rpmesp.2022.392.11906>.
- [3] Gullón, P.; Lumbreras, B.; Sánchez-Martínez, F. I.; de Montalvo-Jaaskeläinen, F.; Ruiz-Cantero, M. T. Public Health and Health Administration in the COVID-19 Pandemic. *SESPAS Report 2022. Gac. Sanit.* **2022**, *36*, S1–S3. <https://doi.org/10.1016/j.gaceta.2022.05.002>.
- [4] Alvarez-Risco, A.; Del-Aguila-Arcentales, S.; Yáñez, J. A. Telemedicine in Peru as a Result of the COVID-19 Pandemic: Perspective from a Country with Limited Internet Access. *Am. J. Trop. Med. Hyg.* **2021**, *105* (1), 6–11. <https://doi.org/10.4269/ajtmh.21-0255>.
- [5] Alexander, G. C.; Tajanlangit, M.; Heyward, J.; Mansour, O.; Qato, D. M.; Stafford, R. S. Use and Content of Primary Care Office-Based vs Telemedicine Care Visits during the COVID-19 Pandemic in the US. *JAMA Netw. Open* **2020**, *3* (10), 1–11. <https://doi.org/10.1001/jamanetworkopen.2020.21476>.
- [6] Del Llano Núñez-Cortés, A. Derecho de La Salud Pública: ¿hacia Un Marco Jurídico de Pandemias? *Gac. Sanit.* **2023**, *37*, 1–2. <https://doi.org/10.1016/j.gaceta.2022.102262>.
- [7] Silva, C. R. D. V.; Lopes, R. H.; Júnior, O. D. G. B.; Fuentealba-Torres, M.; Arcêncio, R. A.; Da Costa Uchôa, S. A. Telemedicine in Primary Healthcare for the Quality of Care in Times of COVID-19: A Scoping Review Protocol. *BMJ Open* **2021**, *11* (7), 1–5. <https://doi.org/10.1136/bmjopen-2020-046227>.
- [8] Equipo técnico de la Dirección de Telesalud, R. y U. *Reglamento de La Ley N°30421, Ley Marco de Telesalud y Del Decreto Legislativo Que Fortalece Los Alcances de La Telesalud.*; Peru, 2021. <http://bvs.minsa.gob.pe/local/MINSA/5418.pdf>.
- [9] OPS. *ESTRATEGIA Y PLAN DE ACCION SOBRE LA PROMOCION DE LA SALUD EN EL*




- CONTEXTO DE LOS OBJETIVOS EN EL CONTEXTO DE LOS OBJETIVOS DE DESARROLLO SOSTENIBLE 2019-2030*; 2019.
- [10] Bashshur, R. L.; Howell, J. D.; Krupinski, E. A.; Harms, K. M.; Bashshur, N.; Doarn, C. R. The Empirical Foundations of Telemedicine Interventions in Primary Care. *Telemed. e-Health* **2016**, *22* (5), 342–375. <https://doi.org/10.1089/tmj.2016.0045>.
- [11] Diaz-Lazo, A.; Montalvo Otivo, R.; Lazarte Nuñez, E.; Aquino Lopez, É.; Montalvo, J.; Diaz-Meyzan, L. Factores Asociados a Mortalidad En Pacientes Con Covid-19 En Un Hospital Público. *Boletín Malariol. y Salud Ambient.* **2022**, *62* (2), 233–240. <https://doi.org/10.52808/bmsa.7e6.622.014>.
- [12] Barcelona Health, H. Estudio de La Consulta de Salud Virtual Telemedicina y Sus Beneficios Para Los Sistemas Sanitarios. **2020**, 1–78.
- [13] Eberly, L. A.; Kallan, M. J.; Julien, H. M.; Haynes, N.; Khatana, S. A. M.; Nathan, A. S.; Snider, C. Patient Characteristics Associated With Telemedicine Access for Primary and Specialty Ambulatory Care During the COVID-19 Pandemic. *JAMA Netw.* **2020**, *3* (12), 1–12. <https://doi.org/10.1001/jamanetworkopen.2020.31640>.
- [14] Orozco-Beltran, D.; Sánchez-Molla, M.; Sanchez, J. J.; Mira, J. J. Telemedicine in Primary Care for Patients with Chronic Conditions: The Valcrònic Quasi-Experimental Study. *J. Med. Internet Res.* **2017**, *19* (12), 1–13. <https://doi.org/10.2196/jmir.7677>.
- [15] Andrews, E.; Berghofer, K.; Long, J.; Prescott, A.; Caboral-Stevens, M. Satisfaction with the Use of Telehealth during COVID-19: An Integrative Review. *Int. J. Nurs. Stud. Adv.* **2020**, *2*, 1–9. <https://doi.org/10.1016/j.ijnsa.2020.100008>.
- [16] Montenegro, P.; Pinillos, L.; Young, F.; Aguilar, A.; Tirado-Hurtado, I.; Pinto, J. A.; Vallejos, C. Telemedicine and the Current Opportunities for the Management of Oncological Patients in Peru in the Context of COVID-19 Pandemic. *Crit. Rev. Oncol. Hematol.* **2021**, *157*, 1–5. <https://doi.org/10.1016/j.critrevonc.2020.103129>.
- [17] Itamura, K.; Rimell, F. L.; Illing, E. A.; Higgins, T. S.; Ting, J. Y.; Lee, M. K.; Wu, A. W. Of Patient Experiences in Otolaryngology Assessment Virtual Visits During the COVID-19 Pandemic. *OTO Open* **2020**, *4* (2), 19–22. <https://doi.org/10.1177/2473974X20933573>.
- [18] Marques, S.; Cruz, J. A. W.; da Cunha, M. A. V. C.; Tuon, F. F.; de Moraes, T. P.; Daiane Zdziarski, A.; Bomher, S. T.; Donnelly, L. F.; Capasso, R. Patient and Family Experience with Telemedicine and In-Person Pediatric and Obstetric Ambulatory Encounters throughout 2020, during the COVID-19 Epidemic: The Distance Effect. *BMC Health Serv. Res.* **2022**, *22* (1), 1–8. <https://doi.org/10.1186/s12913-022-08037-8>.
- [19] Scofano, R.; Monteiro, A.; Motta, L. Evaluation of the Experience with the Use of Telemedicine in a Home Dialysis Program—a Qualitative and Quantitative Study. *BMC Nephrol.* **2022**, *23* (1), 1–9. <https://doi.org/10.1186/s12882-022-02824-5>.
- [20] Frydman, J. L.; Li, W.; Gelfman, L. P.; Liu, B. Telemedicine Uptake among Older Adults during the COVID-19 Pandemic. *Ann Intern Med* **2022**, *175* (1), 145–148. <https://doi.org/10.7326/M21-2972>. Telemedicine.
- [21] Ryskina, K. L.; Shultz, K.; Zhou, Y.; Lautenbach, G.; Brown, R. T. Older Adults ' Access to Primary Care : Gender , Racial , and Ethnic Disparities in Telemedicine. *J. Am. Geriatr. Soc.* **2021**, No. June, 1–9. <https://doi.org/10.1111/jgs.17354>.
- [22] Lutz de Araujo, A.; Moreira, T. de C.; Rados, D. R. V.; Gross, P. B.; Molina-Bastos, C. G.; Katz, N.; Hauser, L. The Use of Telemedicine to Support Brazilian Primary Care Physicians in Managing Eye Conditions : The TeleOftalmo Project. *PLoS One* **2020**, *15* (4), 1–12. <https://doi.org/10.1371/journal.pone.0231034>.
- [23] Holcomb, D.; Faucher, M. A.; Bouzid, J.; Quint-Bouzid, M.; Nelson, D. B.; Duryea, E. Patient Perspectives on Audio-Only Virtual Prenatal Visits amidst the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Pandemic. *Obstet.*





- Gynecol.* **2020**, *136* (2), 317–322. <https://doi.org/10.1097/AOG.0000000000004026>.
- [24] Yadav, D.; Bhatia, S.; Ramam, M.; Singh, V.; Khanna, N.; Khandpur, S.; Gupta, V. Patient Perception and Satisfaction with a Smartphone-Based Teledermatology Service Initiated during the COVID-19 Pandemic at a Tertiary Care Hospital in North India. *Indian J. Dermatol. Venereol. Leprol.* **2022**, *88* (5), 623–632. [https://doi.org/10.25259/IJDVL\\_608\\_2021](https://doi.org/10.25259/IJDVL_608_2021).
- [25] Polinski, J. M.; Barker, T.; Gagliano, N.; Sussman, A.; Brennan, T. A.; Shrank, W. H. Patients' Satisfaction with and Preference for Telehealth Visits. *J. Gen. Intern. Med.* **2016**, *31* (3), 269–275. <https://doi.org/10.1007/s11606-015-3489-x>.
- [26] Contractor, U.; Hass, W.; Reed, P.; Osborne, L.; Tree, J.; Bosanquet, D. C. Patient Satisfaction with Tele- and Video-Consultation in the COVID-19 Era e A Survey of Vascular Surgical Patients. *Ann VascSurg* **2020**, *85*, 105–109. <https://doi.org/https://doi.org/10.1016/j.avsg.2022.05.009>.
- [27] Carrillo de Albornoz, S.; Sia, K. L.; Harris, A. The Effectiveness of Teleconsultations in Primary Care: Systematic Review. *Fam. Pract.* **2022**, *39* (1), 168–182. <https://doi.org/10.1093/fampra/cmab077>.
- [28] Rodriguez, H. El Impacto de La Teleconsulta En Los Derechos Fundamentales de La Población Aislada – Rural, Conforme Reglamento de La Ley N°30421, Universidad cesar Vallejo, 2021. <https://hdl.handle.net/20.500.12692/83825>.
- [29] Harrigan, M.; Cartmel, B.; Loftfield, E.; Sanft, T.; Chagpar, A. B.; Zhou, Y.; Playdon, M.; Li, F.; Irwin, M. L. Randomized Trial Comparing Telephone versus In-Person Weight Loss Counseling on Body Composition and Circulating Biomarkers in Women Treated for Breast Cancer: The Lifestyle, Exercise, and Nutrition (LEAN) Study. *J. Clin. Oncol.* **2016**, *34* (7), 669–676. <https://doi.org/10.1200/JCO.2015.61.6375>.
- [30] Lindegaard Pedersen, J.; Pedersen, P. U.; Damsgaard, E. M. Nutritional Follow-up after Discharge Prevents Readmission to Hospital - A Randomized Clinical Trial. *J. Nutr. Heal. Aging* **2017**, *21* (1), 75–82. <https://doi.org/10.1007/s12603-016-0745-7>.
- [31] Galan, G. Calidad de Atención y Satisfacción de Los Pacientes Atendidos Por Telemonitoreo y Teleorientación En La Posta Médica Túcume – EsSalud, Universidad Cesar Vallejo, 2021. <https://hdl.handle.net/20.500.12692/81104>.
- [32] Mizziara, R. A.; Maesaka, J. Y.; Matsumoto, D. R. M.; Pentead, L.; Anacleto, A. A. D. S.; Accorsi, T. A. D.; Lima, K. D. A.; Cordioli, E.; D'alessandro, G. S. Teleoncology Orientation of Low-Income Breast Cancer Patients during the COVID-19 Pandemic: Feasibility and Patient Satisfaction. *Rev. Bras. Ginecol. e Obstet.* **2021**, *43* (11), 840–846. <https://doi.org/10.1055/s-0041-1739425>.
- [33] Thomas, N. A.; Drewry, A.; Racine Passmore, S.; Assad, N.; Hoppe, K. K. Patient Perceptions, Opinions and Satisfaction of Telehealth with Remote Blood Pressure Monitoring Postpartum. *BMC Pregnancy Childbirth* **2021**, *21* (1), 1–11. <https://doi.org/10.1186/s12884-021-03632-9>.
- [34] Kirkland, E.; Schumann, S. O.; Schreiner, A.; Heincelman, M.; Zhang, J.; Marsden, J.; Mauldin, P.; Moran, W. P. Patient Demographics and Clinic Type Are Associated with Patient Engagement within a Remote Monitoring Program. *Telemed. e-Health* **2021**, *27* (8), 843–850. <https://doi.org/10.1089/tmj.2020.0535>.
- [35] Kirkland, E. B.; Marsden, J.; Zhang, J.; Schumann, S. O.; Bian, J.; Mauldin, P.; Moran, W. P. Remote Patient Monitoring Sustains Reductions of Hemoglobin A1c in Underserved Patients to 12 Months. *Prim. Care Diabetes* **2021**, *15* (3), 459–463. <https://doi.org/10.1016/j.pcd.2021.01.005>.
- [36] Randal, M. H.; Haulsee, Z. M.; Zhang, J.; Marsden, J.; Moran, W. P.; Kirkland, E. B. The Effect of Remote Patient Monitoring on the Primary Care Clinic Visit Frequency among Adults with Type 2 Diabetes. *Int J Med Inf.* **2020**, *143* (5), 139–148. <https://doi.org/10.1016/j.ijmedinf.2020.104267>.
- [37] Stulock, R.; Montgomery, J.; Parker, M.; Soric, A.; Zeleznikar, E. Pharmacist Involvement in a



- Comprehensive Remote Monitoring and Telemanagement Program. *Am. J. Heal. Pharm.* **2022**, 79 (11), 888–895. <https://doi.org/10.1093/ajhp/zxac025>.
- [38] Hanley, J.; Pinnock, H.; Paterson, M.; McKinsty, B. Implementing Telemonitoring in Primary Care: Learning from a Large Qualitative Dataset Gathered during a Series of Studies. *BMC Fam. Pract.* **2018**, 19 (1), 1–11. <https://doi.org/10.1186/s12875-018-0814-6>.
- [39] Sin, D. Y. E.; Guo, X.; Yong, D. W. W.; Qiu, T. Y.; Moey, P. K. S.; Falk, M. R.; Tan, N. C. Assessment of Willingness to Tele-Monitoring Interventions in Patients with Type 2 Diabetes and/or Hypertension in the Public Primary Healthcare Setting. *BMC Med. Inform. Decis. Mak.* **2020**, 20 (1), 1–11. <https://doi.org/10.1186/s12911-020-1024-4>.
- [40] Georghiou, T.; Sherlaw-Johnson, C.; Massou, E.; Morris, S.; Crellin, N. E.; Herlitz, L.; Sidhu, M. S.; Tomini, S. M.; Vindrola-Padros, C.; Walton, H.; Fulop, N. J. The Impact of Post-Hospital Remote Monitoring of COVID-19 Patients Using Pulse Oximetry: A National Observational Study Using Hospital Activity Data. *eClinicalMedicine* **2022**, 48, 101441. <https://doi.org/10.1016/j.eclinm.2022.101441>.
- [41] Beaney, T.; Clarke, J.; Alboksmaty, A.; Flott, K.; Fowler, A.; Bengler, J.; Aylin, P. P.; Elkin, S.; Neves, A. L.; Darzi, A. Population-Level Impact of a Pulse Oximetry Remote Monitoring Programme on Mortality and Healthcare Utilisation in the People with COVID-19 in England: A National Analysis Using a Stepped Wedge Design. *Emerg. Med. J.* **2022**, 39 (8), 575–582. <https://doi.org/10.1136/emermed-2022-212378>.
- [42] Gutierrez, B. G. CALIDAD DE LAS TELEINTERCONSULTAS EN LA RED ASISTENCIAL DE MOQUEGUA 2021 PRESENTADO POR SERVICIOS DE SALUD LIMA – PERÚ, Universidad San Martín de Porres, Perú, 2022. <https://hdl.handle.net/20.500.12727/10615>.
- [43] Tostes, J. G.; Benedita, C.; Cruz, R. P.; Mota, G.; Assis, C. De; Alves, R. Tele-Interconsultation and the Challenges for Maintaining Health Care in Pandemic Times. *Rev Cienc Saude* **2021**, 11 (2), 5–9. <https://doi.org/10.21876/rchsci.v11i2.1069>.