Journal of Chemical Health Risks

www.jchr.org

JCHR (2023) 13(5), 883-886 | ISSN:2251-6727



Assessing the Patient Profiles of Individuals with Diabetic Foot Conditions

Prasidh Vijay Shetty(Junior resident)^{1,*}, S. J. Bhosale(Professor)¹ and Saurabh Tukaram Gondil(Senior resident)¹

¹Department of General Surgery, Krishna Institute Medical Sciences, KVV, Karad, Maharashtra, India.

Corresponding author: Prasidh Vijay Shetty (Junior resident)

Department of General Surgery, Krishna Institute Medical Sciences, KVV, Karad, Maharashtra, India.

(Received: 28	October 2023	Revised: 08 November	Accepted: 02 December)
KEYWORDS	ABSTRACT:		
antibiotic sensitivity,	The objective o	f the research was to analyze both th	e clinical and bacterial characteristics of
anaerobic,	individuals suffe	ring from diabetic foot conditions. Th	is comprehensive investigation spanned a
limb	two-year period	and involved patients diagnosed with D	iabetes mellitus. During the study, the foot
	ulcers were class	ified into two categories: mild, which we	ere non-limb threatening, and severe, which
	posed a threat to	b limb integrity. Prior to the initial deb	pridement procedure, wound cultures were
	collected, and an	ntibiotic sensitivity testing was conduc	ted for each case. Additionally, the study
	documented pote	ential risk factors contributing to the de	evelopment of these ulcers. Among the 60
	cases of diabetic	e foot ulcers, 36 patients (62.29%) exp	perienced non-limb threatening infections,
	while 24 patients	s (37.7%) faced limb-threatening infect	ions. Within this group of 24 patients with
	limb-threatening	infections, 16 individuals (60.87%) ultit	mately required amputation during the one-
	year follow-up p	eriod. The microorganisms identified from	m the cultures encompassed various strains,
	with Staphyloco	ccus species appearing in 14 patients (28	3.84%), Streptococcus species in 10 patients
	(21.15%), Pseudo	omonas species in 6 patients (15.38%), En	nterococcus species in 20 patients (57.69%),
	and anaerobic ba	cteria in 10 patients (51.92%).	

I. INTRODUCTION

Diabetes mellitus has emerged as an epidemic in the Asian sub-continent, particularly impacting India where nearly 20 million individuals are affected. Among the complications of diabetes, diabetic foot ulcers are a significant concern, esti- mated to afflict approximately 15% of all diabetics over their lifetime. Alarmingly, these ulcers precede almost 80% of foot amputations [1], [2]. The complex interplay of diabetes- related factors such as neuropathy, vasculopathy, and altered musculoskeletal and soft tissue mechanics contributes to elevated plantar pressure. This heightened pressure predis- poses individuals to tissue damage, particularly at weight- bearing sites, culminating in non-resolving neuro-ischemic ulcers. Notably, a majority of skin injuries in diabetic patients manifest on the plantar surface, often occurring at sites ex- periencing the highest pressure beneath the foot [3], [4]. The significance of this current study becomes particularly

pro- nounced within the Indian context. Here, the disease is often diagnosed at an advanced stage, awareness of proper foot care among patients is limited, and there is a substantial delay in seeking medical treatment. Moreover, a considerable portion of the population resides in rural areas and engages in fieldwork without adequate footwear, which significantly heightens the risk of additional infections. In this challenging scenario, physicians are often compelled to initiate empirical treatment while awaiting culture reports. Therefore, having a preliminary understanding of the antibiotic sensitivity pat- terns proves to be a valuable tool for healthcare providers in their decision-making process.

II. METHODS

The study encompassed a cohort of 60 patients diagnosed with diabetic foot conditions, comprising both hospitalized individuals and outpatients. Prior to

Journal of Chemical Health Risks www.jchr.org

JCHR (2023) 13(5), 883-886 | ISSN:2251-6727



their inclusion in the study, each patient provided informed consent, and the study protocol adhered to the ethical principles outlined in the 1970 Declaration of Helsinki. The average age of these patients stood at 50.4 years, spanning an age range of 30 to 66 years. The study group consisted of 32 male participants and 28 female participants.

The classification of foot ulceration in this study was based on severity, distinguishing between mild or nonlimb threatening and severe or limb-threatening ulcers. Non-limb threatening ulcers were defined as superficial ulcers with less than 2cm of cellulitis, no signif- icant ischemia, and no bone or joint involvement, and

without systemic toxicity4. On the other hand, limbthreatening ulcers were described as full-thickness ulcers with more than 2cm of cellulitis, severe ischemia or gan- grene, bone or joint involvement, and associated systemic toxicity. Systemic toxicity was determined by the pres- ence of clinical symptoms along with elevated markers- ESR, leukocytosis and polymorphonuclearcytosis. The investigations in this study followed the protocol of the diabetic clinic. The tests conducted encompassed random blood sugar, fasting and post-prandial blood sugar levels, Glycosylated hemoglobin assessment, routine and microscopic urine analysis, renal function tests (including BUN and serum creatinine), liver function tests, a chest X-ray (PA view), ECG, and fundus examination. These comprehensive investigations were conducted to assess various aspects of the patients' health relevant to their diabetic condition and foot ulceration. Each case in the study underwent a thorough assessment, including aerobic and anaerobic culture along with antibiotic sensitivity testing. Additionally, a complete blood count with ESR (erythrocyte sedimentation rate) was conducted. To further characterize the ulcers and determine any evidence of bone involvement, a probe-to-bone test was performed. Imaging studies, such as X-ray of the wound and MRI, were conducted for all patients. These comprehensive evaluations aimed to provide a comprehensive understanding of the ulcer's nature and extent, facilitating appropriate treatment decisions.

Organism	No.ofpatients	Percentage
Staphylococcussp.	14	28.84
Streptococcussp	10	21.15
Pseudomonas	6	15.38
Enterococcussp.	20	57.69
Anaerobes	10	51.92

TABLE 1: Base line characteristics of the study group N=60

III. RESULTS

Among the 60 patients included in the study, wound cultures were conducted for all cases. Following an incubation period of 48-72 hours, the microbiological profile obtained from these cultures is presented in Table 2. This profile likely includes information about the types of microorganisms iden tified and their respective proportions in the wound cultures, shedding light on the bacterial composition associated with diabetic foot ulcers.

In an attempt to ascertain the potential factors contribut-

ing to the development of diabetic foot, a comprehensive investigation was carried out. A detailed patient history was collected, with a focus on relevant factors. Additionally, a thorough examination of the foot was conducted to identify pressure points on the sole. Through these assessments, ef- forts were made to uncover and understand the factors that might be associated with the onset or progression of diabetic foot complications.

www.jchr.org

JCHR (2023) 13(6), 1334-1342 | ISSN:2251-6727



Antibiotic	No. of patients	Percentage
Amoxycillin-clavulanicacid	57	94.23
Ceftrioxone	56	92.30
Ceftazidime	49	80.32
Cefuroxime	42	66.85
Cephalexin	33	52.09
Cephadroxil	31	50.81
Ampicillin	34	55.73
Cloxacillin	44	70.13
Co-trimexazole	46	74.40
Ciprofloxacillin	52	86.24
Ofloxacillin	51	83.60
Sparfloxacillin	51	81.60
Chloremphenicol	55	90.16
Nitrofurantoin	52	84.24
PolymyxinB	53	89.88
Gentamycin	47	77.04
Amikacin	55	90.16

TABLE 2: Microbiological profile of patients N=60

TABLE 3: Anti bio ticsensitivity pattern of patients N=60

Parameters		
Age(years)	50.4+/-11.77years	
Range	30-66years	
Sex	32	
MalesFemales	28	
Mean duration of diabetes Foot ulcer	8.6+/-3.2years	
Non-limb threatening	36 (62.29%)	
Limb threatening	24(37.7%)	

IV. DISCUSSION

India is home to a staggering population of approximately 20 million individuals with diabetes, making it the highest in the world. Alarmingly, nearly 15% of these individuals suffer from the serious complications associated with diabetic foot [5]. Diabetics in India face a unique set of challenges due to sociocultural practices, such as walking barefoot, engaging in religious practices like walking on fire, using inadequate footwear, and a lack of awareness regarding proper foot care. These factors collectively contribute to the heightened preva- lence of diabetic foot complications in the country, presenting a significant healthcare concern. Diabetic foot ulcers don't develop spontaneously but result from a complex interplay of various factors. These include neuropathy, autonomic neu- ropathy, peripheral vascular disease, along with changes in plantar pressure, inadequate footwear, and restricted joint mobility. Furthermore, cell-mediated immunity is particu- larly affected, leading to abnormalities in key immune cells such as polymorphonuclear leukocytes (PMNL), monocytes, and lymphocytes. These abnormalities manifest in impaired functions like adherence, chemotaxis, phagocytosis, oxida- tive burst, and intracellular killing, collectively contributing to the development and exacerbation of diabetic foot ulcers. In addition to the noted abnormalities in innate immunity, adaptive cellular immunity is also impacted, leading to a reduced lymphocyte proliferative response to stimuli like Staphylococcus aureus. In the current study, the isolation of S. aureus was observed in nearly 14% of

Journal of Chemical Health Risks

www.jchr.org

JCHR (2023) 13(6), 1334-1342 | ISSN:2251-6727



patients, a significant finding. The culture yield was enhanced by ob- taining deep wound samples, which likely contributed to this observation. Interestingly, similar patterns of microbial growth have been reported in other studies, such as the one conducted by Lipsky et al [1]. However, in our study, Enterococcus species and anaerobic bacteria were the most prevalent among the isolated microorganisms. In our study, it was found that ill-fitting shoes and shoerelated condi- tions were more common contributors to diabetic foot issues than the habit of walking barefoot. This prevalence may be attributed to the hilly terrain, economic challenges, and the improper distribution of pressure on the sole associated with traditional footwear. Additionally, the relatively cooler climate in the region discourages barefoot walking. Trivial trauma and ingrown nails accounted for approximately 20% of these cases. Interestingly, in 5 cases (8.19%), no specific cause for diabetic foot issues could be definitively identified.

V. CONCLUSION

Enterococcus species, Staphylococcus species, and anaer- obic bacteria were identified as the predominant culprits behind the majority of foot ulcers in this study. Importantly, these microorganisms displayed sensitivity to conventional antibiotics, suggesting no significant evidence of antibiotic resistance. Therefore, the empirical use of these antibiotics, whether administered orally or by injection (depending on the type of foot ulcer), is justified. Additionally, it is strongly recommended to provide proper education to patients on the importance of appropriate footwear and foot care to prevent and manage diabetic foot ulcers effectively [6], [7].

FUNDING

This research did not receive any specific grant from funding agencies in the public, commercial, or nonprofit sectors.

CONFLICTS OF INTEREST

The authors declared no conflict of interest.

AUTHORS' CONTRIBUTIONS

All authors equally contributed to preparing this article.

References

- Lipsky BA. Evidence based antibiotic therapy of diabetic foot infections. FEMS Immunology and Medical Microbiology 1999; 26: 267.
- [2] Deresinski S. Infections in diabetic patients: Strategies for clinicians. Infectious Disease Reports 1995; 1:1.
- [3] Lipsky BA. Osteomyelitis of the foot in diabetic patients.Clin Infect Dis 1997; 25:1318.
- [4] Pendsey SP. Epidemological aspects of diabetic foot.Int J Diab Dev Countries 1994; 14: 37-38
- [5] Boultrom AJM. The pathway to ulceration: Aetiopathogenesis.BoultonAJM,ConnerH,Cavanagh PR (eds). The Foot in Diabetes.3rd edition. John Willy and sons Ltd, Chichester, West Sussex: 2000 ; pp 19-31.
- [6] Reiber GE, Vileikyte L, Boyke EJ. Asual pathways for incident lower extremity ulcers in patients with diabetes from two settings. Diabetes Care 1999; 22 : 157-162.
- [7] Caputo GM, Cavanagh PR, Ulbrecht JS et al. Current concepts: Assess- ment and management of foot disease in patients with diabetes. N Eng J Med 1994; 331: 854.