



Quality by Design Assisted Revolutionizing Canagliflozine Analysis and Validation with Green Chemistry Innovations

Diptimayee Jena¹, Sivasankar Mohanty², Pratikshya Sarangi³, Shruti Jha³, Jeevika Kumari³, Kirtimaya Mishra^{1*}

^{1,1*} Centurion University of Technology and Management, Bhubaneswar, Odisha

² Seemanta Institute of Pharmaceutical Sciences, Jharpokharia, Mayurbhanj, Odisha

³ ARKA JAIN University, Jamshedpur, Jharkhand

(Received: 07 January 2024

Revised: 12 February 2024

Accepted: 06 March 2024)

KEYWORDS

Green Chemistry, Validation, Eco-friendly, Spectrophotometric, Canagliflozine, Quality by design.

ABSTRACT:

Introduction: Green chemistry, typically referred to as ecologically sustainable chemistry, constitutes a field of inquiry situated within the context of chemistry, wherein the primary objective is to mitigate the adverse impacts related to chemical processes and products.

Objectives: The objective is to promote the advancement of sustainable and eco-friendly treatments by decreasing or completely removing the utilisation of toxic chemicals, minimising the production of hazardous materials, and saving energy and resources. This study observed linearity with a range of concentration between 02-12 μ g/mL for Canagliflozin (CGFZ).

Methods: The method of solving concurrent equations was utilised to ascertain the concentrations of the medication. The recovery percentage from the parenteral dosage form was determined to be 100.4% along with a standard deviation (SD) of ± 0.028 , with a sample size of 6. The recovery rates observed in the accuracy study ranged from 99.8-101.05% on average. The observed percentage relative standard deviation (%RSD) demonstrated a value that was much below 2% throughout intra-day evaluations, suggesting a noteworthy degree of precision in the proposed methodology.

Results: The findings from statistical analyses of the technique the validation outcomes indicate that the suggested procedures are suitable for implementation inside the facilitated quality control laboratories.

Conclusions: The present methodology is deemed appropriate for the quantitative analysis of CGFZ in parenteral dosage formulations, since it effectively eliminates any potential interference from frequently used additives. Hence, this approach can be employed for regular analytical objectives.

1. Introduction

The entire globe is now experiencing revenue generation as a result of the increased advancement of technological and scientific developments, but this economic growth also contributes to ecosystematical degradation, which is demonstrated by issues like climate change, ozone holes, and the buildup of non-invasive organic contaminants throughout the natural world [1]. To reduce the negative effects on the natural world resulting from the manufacture and dissemination of chemicals, a multidisciplinary approach known as "green chemistry"

has been settled. Regarding the sake of reducing waste, saving assets, and increasing the use of organic items, the optimum focus of this study is the development of environmentally friendly alternatives that substitute traditional procedures utilising chemicals [2]. Canagliflozine (CGFZ), denoted by its IUPAC (International Union of Pure and Applied Chemistry) symbol (2S,3R,4R,5S,6R)-2-{3-[5-[4-Fluoro-phenyl]-thiophen-2-ylmethyl]-4-methyl-phenyl}-6-hydroxymethyl-tetrahydro-pyran-3,4,5-triol [3]. By preventing the kidneys from reabsorbing glucose, CGFZ lowers blood glucose levels by increased amount of



glucose excreted in the urine [4]. CGFZ, which has been approved by regulatory bodies including the FDA and the EMA, is now a crucial part of all-encompassing treatment plans for individuals with type 2 diabetes, providing alternatives for both monotherapy and combination therapy [5].




Figure 1. Chemical structure of CGFZ

CGFZ in injectable forms & biological specimen is estimated throughout UV-Visible Spectroscopy, HPLC along with LC-MS method [6]. The disclosed UV spectrophotometric approach, however, has a number of limitations. These include a limited linearity range, no Sandell's sensitivity, and no presentation of the molar extinction coefficient. As a result Quality-by-Design (QbD) strategy was implemented to the advancement of a novel as well as developed UV spectrophotometric method for quantification of CGFZ through tablet preparation [7].

A holistic strategy, QbD, that incorporates excellence throughout every aspect of the operation to guarantee the desired outcome is achieved. QbD is a framework for systematic creativity that conforms, to the standards set out by International Council for Harmonisation (ICH) of Technical Requirements, for Pharmaceuticals for Human Use, in particular ICH-Q8-(R2). This approach begins with clearly defined objectives and places a strong focus on comprehending and maintaining both the final outcome and its structure. This strategy's technique is based on good scientific concepts and incorporates measures that lower risks, guaranteeing the results' accuracy and dependability. After Food and Drug Administration (FDA) launched its "Pharmaceutical Current Good Manufacturing Practises (cGMPs) for the 21st Century" in 2002, QbD has been launched. There are six stages involved in implementing the empirical QbD Perspective and achieving a dependable and high-quality analysis approach with increased resilience.

The implementation of the QbD methodology not only decreases the time required for developing a reliable analytical technique but is also considered financially

efficient strategy to ensuring quality from the early stages of method development. For determining the optimum configuration domain for method efficiency, making it an integral part of paradigm, QbD, Design of Experiments (DoE), is widely regarded like a crucial tool. The use of rigorous experimental designs is the primary focus of this investigation; the overarching goal is to reduce the amount of variation that occurs in spectrophotometric aspect of CGFZ. Primary objective is to ascertain the finest selections. The investigation commenced with the implementation of a factor screening analysis utilising a fractional factorial design (FFD) in order to ascertain the key method parameters that exert an influence on performance. Subsequently, a central composite design (CCD), was employed to optimise the methodology, so ensuring its resilience and attaining predetermined objectives. The goal of the study was to come up with an innovative UV spectrophotometric method that was rigorous and reliable for determining the amount of CGFZ in tablet formulations. We were able to accomplish this objective thanks to the effective application of QbD concepts throughout the method development process, followed by validation in accordance with ICH standards.

2. Materials and Methods

Reagents and Standards

CGFZ standardised sample with purity >99.5% was provided by Dolphin Pharmaceutical, Surat, India. Methanol (MeOH) utilised in the preparation of our medication and reagent solutions came from Merck Ltd., Jamshedpur, India. As the marketed tablet formulation of CGFZ (100mg) was accessible, it was purchased and analysed using the established procedure.

Instrumentation and Optical Characteristics

The spectrum investigations were carried out via a microprocessor-controlled single beam system and the LI-285 UV spectrophotometer (manufactured by Lasany, India), employing 10mm calibrated quartz cuvettes. To ensure precise measurements of reagents, a high-precision analytical instrument was employed. Ultrasonication (Enertech., India), was utilized to influence the dissolution of selected dosage form.

Incorporation of Analytical Target Profile

It's one of the comprehensive examinations of existing literary text sources & medication profiles, including physical as well as chemical features, were conducted to



develop an analytical target profile. This profile encompasses a concise overview of the quality attributes associated with an analytical method. Ultimate focus of this study, was to create an analytical method that is efficient, accurate, and economical for estimating the concentration of CGFZ in its tablet dose form. Hence, in accordance with the primary objective of this research, a UV spectrophotometric technique was employed to facilitate the expeditious analysis of CGFZ. The decision to employ the UV spectrophotometric method in drug analysis was motivated by its advantageous characteristics of simplicity and efficiency, which render it superior to more complex analytical techniques.

Incorporation of Risk Management & Cause-Effect Relationship

When it's important to see how different factors that might affect the performance of a method are connected, the Ishikawa fish-bone diagram is one of the easiest ways to do it. An Ishikawa diagram was made to learn more about how these factors might change the UV spectrophotometry properties for CGFZ. Researchers used Cause-Effect Risk Assessment Matrixes depending on the Control-Noise-Experimentation (CNX) method to figure out which factors are most likely to affect the study's statistical features. The study found lots of CMVs i.e., Critical Method Variables that were bridged to elevated end score, which suggests that they are high-risk variables. The composition of those CMVs encompasses variations in detection wavelength, solvent used, scan rate, sample integrity, sampling intensity, and pH of the sample. The CMVs were checked out by implementing screening design additionally in order to determine the crucial method parameters. Subsequently, response surface optimisation was conducted using an appropriate experimental design.

Screening of CMVs by FFD

To identify the variables with high risk, Design expert 13 software, version-13.0.14, USA, FFD has been utilized to screen critical variables. Several factors were chosen as essential method variables based on a comparison of the spectrum design, precision, and absorbance. Based on what was already known and the Ishikawa fish-bone diagram, prioritisation studies were used to figure out the detection wavelength, solvent type, & sample integrity. These factors further checked using direct observation. Design expert software was used to conduct an FFD experiment with a minimum of 5 trials (one serving as a

centre point) to assess the technique variables of scan speed (X1), PH of the sample (X2), and interval of sampling (X3). Parameters were evaluated on both the higher and lower levels, as well as the program was further utilized in determining CMVs those affect the absorbance (Y) of the response variables. Significant parameters were identified by evaluating actual versus predicted values plot, prediction equation, pareto chart & fitting summary plot.

Method optimization & Robustness study Implementing CCD

The CCD was used to make sure that the method for finding the best method conditions was reliable. As a result of the screening studies, 13 dummy runs were ready using at the least five central points chosen rely on CCD to find the best CMVs, such as the Conc. of MeOH (A) & pH (B). As the answer variable, absorbance at 290 nm was used to look at the results of the experiments. For all the tests, a normal CGFZ of 10µg/mL was used.

Design of Expert software was used to fit experiment data into mathematical model applying Multiple Linear Regression Analysis (MLRA). This model could explore main and interaction effects. To analyse the model, only significant coefficients (p value<0.05) from analysis like ANOVA were used for framing the polynomial equations & analyse parameters for instance R², adjusted R², & Predicted Residual Sum of Squares (PRESS) respectively. Multiple profilers were used to evaluate the model's viability, including projection profilers, interaction profilers, and three-dimensional response surface profilers. Using a numerical desirability function, we were able to find the optimal solution by striking a balance between the many factors under consideration. The area of the design was then demarcated off according to this.

Method Control Strategy

The design space formed by the DoE methodology was used to inform the development of control strategies for the method, enclosed by the marginally changes in method performances were enabled to retain method's resilience.

Preparation of Standard Stock Solution

To make the CGFZ standard stock solution (1000µg/mL), 10 mg of CGFZ was dissolved in enough EtOH to make 10 mL. To make 100 g/mL standard solutions, five mL of above prepared stock solution



transferred, to a fifty mL volumetric flask, & the content was filled with water to make fifty mL.

Parenteral Dosage Form Analysis

Invokanna 100/Jassen Pharmaceutical labelling claim for its CGFZ tablet formulation is 100 mg. For tablet, 10 mg equivalent weight of CGFZ should be diluted in 10 millilitres of MeOH. After the solution was ready, 5ml of solution was transferred to a 50 millilitre volumetric flask and brought to 50 millilitres with water. For thirty minutes, the material was ultrasonicated. Whattmann filter paper was used to further filter this solution for particles. In order to analyse the filtered solution, it was diluted further with water. Using a calibration curve based on the concentration of standard CGFZ, we were able to calculate the amount of medication in the sample solution [8].

3. Method Validation

The Specificity

The specificity of UV spectrophotometric method, was appraised by determining drug's presence in the formulation excipients. The spectrums were assessed in order to see if there was any potential interference caused by the excipients [9].

Linearity

In order to create a range of concentrations from 2 to 12 g/mL, multiple aliquots were collected with working standard solution of CGFZ and placed within distinct 10 mL volumetric flasks before diluting by using H₂O at the end of the process [10]. At a wavelength of 290 nm, UV absorption was measured. For the purpose of determining whether or not the results are linear, a calibration curve was created by placing the absorbance

& the concentration (in g/mL) on the Y-axis & X-axis respectively.

Precision & Accuracy

In order to assessing the methodology precision, recovery experiments were being conducted with three different levels: 80%, 100%, & 120% of the test solution of CGFZ (10 μ g/mL), utilising the standard dilution technique. Recovery researches were conducted in duplicate for every degree. Standard medication, CGFZ mixed up with the recovery solution was computed by plotted calibration curve. In order to evaluate intra-day precision, replicates of total six of a constant conc. of CGFZ (10 μ g/mL) were reviewed within a single day, as well as subsequently, the %RSD values were computed.

4. Results

In this current research, Ultraviolet spectrophotometric (UV Spectrophotometry) method was established for the purpose of determining the amount of CGFZ in a tablet formulation. In order to obtain the variable parameters that would be needed for constructing the final spectrophotometric settings, the QbD method was put into practise. For the purpose of determining the technique variables, a conventional Ishikawa fish-bone diagram was constructed. The physical evaluation of the variables involved in the procedure was carried out. It was discovered that the medication did not dissolve in either acetone or ether. CGFZ was soluble in water. The standard CGFZ solution reaches its maximum absorption (λ_{max}) in water at a wavelength of 290 nm (Fig. 2); this wavelength was chosen as the detecting wavelength




Figure 2. Typical UV absorption spectrum of CGFZ



Melting point test showed good sample integrity [11]. However, sample PH, scan speed, sampling interval required for a thorough research for assessing their effects on technique robustness. Screening CMVs by scan speed, conc. of MeOH, & sample PH was made easier using FFD. Actual vs. projected plots showed the model's fitness. Model p-value (0.0005), R2 (0.9335), & RMSE (Root Mean Square Error) (0.0175) indicated

model suitability. Summary of fit showed predicted R2 (0.5269) & adjusted R2 (0.8860).

The CCD was used to check how the CMVs changed the reaction absorbance. A UV spectrophotometer was used for 13 experiments that were done in a random order to get an answer that was free of bias and had at least five centre points. Table 1 shows the results of each experiment and the spectrophotometric range that was looked at.

Table 1. Experimental design matrix showing spectrophotometric range studied for robustness study and obtained responses

Run No.	Conc. of MeOH in ml (A)	pH (B)	Absorbance (Y)
1	4	2	0.388
2	1.58579	4	0.38
3	4.41421	4	0.366
4	2	6	0.468
5	3	4	0.445
6	3	4	0.445
7	3	1.17157	0.316
8	3	4	0.445
9	3	6.82843	0.375
10	4	6	0.382
11	3	4	0.445
12	2	2	0.311
13	3	4	0.445
Range	Low	High	
MeOH	2	4	
pH	2	4	

The null hypothesis (H0) was accepted based on a predetermined significance level of 0.05 for the p-value. A comprehensive examination of the CCD model was conducted, employing various statistical analytic techniques such as ANOVA, parameter estimations, and prediction profiler, to derive meaningful conclusions.

In Fig. 3 (A), perturbation plots display the anticipated models, independent factors impact on a given response, while fixing all other factors constant at a reference point. Degree of steepness in a slope or curve indicates the sensitivity of the reaction to a certain component [12]. In the presented analysis in Fig. 3 (A), it was denoted that factor B, the sampling interval exhibited the greatest significant impact upon absorbance, following the sample pH. The actual versus predicted graphic

showcases the baseline model (represented by blue points), revealing that the line obtained for the experimental data falls smoothly within the accuracy interval parameters (Fig. 3 (B)). The observed data exhibited a high degree of similarity to the expected data, this disproves the null hypothesis and proves the model's efficacy in describing data variation [13].

Response surfaces plots against conc. of MeOH & pH are illustrated in (Fig. 4) (Conc. of MeOH is plotted against the pH). The analysis of perturbation plots & response plots, of optimization models revealed that factor had a huge impact on the absorbance of the analytes.

In addition, the analysis of variance (ANOVA) resulted in a P-value that was found to be less than 0.0005, which indicates that the model is suitable for addressing the



variability observed in the data. This finding also suggests that the null hypothesis should be rejected. Additionally, the model's appropriateness was confirmed by the lowest value observed in anticipating PRESS [14].

The review of parametric estimations is vital in estimating the risk of variability from diverse variables. A p-value that is observed to be less than 0.05 indicates the availability of non-zero slope.

Sampling interval \times pH (B2) and conc. of MeOH (A) became the greatest impacting method variable.

$$\text{Absorbance } (Y) = 0.4450 - 0.0036A + 0.0293B - 0.0407AB - 0.0290A^2 - 0.0427B^2$$

Denoted as, A = Conc. of MeOH, B = pH.




Figure 3 (A). Perturbation Plot, **3(B).** Predicted vs. Actual Plot

Table 2 displays the optical properties associated with the spectrophotometric technique. The method that was devised demonstrated both specificity and selectivity, since it was shown that the frequently utilised formulation additives in the selected dosage form did not interfere with the predicted procedure. The medication exhibited linearity, within a range of concentration 2-12 μ g/ml. The conducted regression analysis, on the linearity data demonstrated a satisfactory level of overall goodness of fit. The statistical measures, namely R², adjusted R², and predicted R², yielded values of 0.9335, 0.8860 and 0.5269, respectively. The analysis of variance (ANOVA) indicated that the approach used for assessing linearity data was deemed adequate, as evidenced by a statistically significant p-value of less than 0.05.

Recovery percentage from the selected dosage form (tablet dose form) was determined to be 100.4% with a standard deviation of ± 0.0175 , based on a sample size of 6.

5. Discussion

The recovery rates observed in the accuracy investigation exhibited an average range of 99.8-101.05%. In relation to intraday assessments, the %RSD exhibited a value well below 2%, indicating a notable level of precision in the suggested methodology. The obtained findings of the employed methodology fall within the predetermined range, indicating that the methodology is unaffected by the presence of additives.




Figure 4. 3-D Response surface plot for absorbance against Conc. of MeOH Vs. pH

Table 2. Optical Characteristics and Summary of validation parameters

Parameters	Obtained Values
Wavelength (nm)	290
Linearity Range ($\mu\text{g/ml}$)	2-12
Molar extinction coefficient ($\text{ltr}/\text{mol.cm}$)	2.00478×10^4
Regression equation ($Y=ax+b$)*	$0.044x + 0.0085$
Correlation coefficient (R^2)	0.9984
Precision (% R.S.D., $n=6$)	0.2092032
Accuracy (% Recovery \pm S.D.)	
80%	0.403707 ± 0.003266
100%	0.782301 ± 0.006944
120%	0.249693 ± 0.002449
% Range of error	
95% confidence limits	± 0.022
99% confidence limits	± 0.030

R.S.D. – Relative Standard Deviation; S.D. – Standard Deviation; A.U. - Absorbance Units,* is $Y = ax + b$, where Y = absorbance, a = slope, b = intercept and x is the concentration, † is average of three determinations at each level

6. Conclusion

An exact UV spectrophotometric approach for measuring CGFZ was made achievable due to the QbD methodology. The QbD procedure guaranteed a high quality of analysis. When establishing control techniques for the method and planning future trials to maintain making the method perform better, the researcher had to pay particular attention to the conc. of MeOH and pH. Based on the data, the method seems to be new, easy, accurate, and precise. The created methods can be used in quality control labs, according to statistical studies of the method validation results. This technique can be used to find out the CGFZ of a tablet dosage form without any problems from widely used fillers. Because of this, particularly this method can be implemented for normal analytical goals.

Acknowledgment

The research team express their gratitude to Jassen Pharmaceutical, India regarding generously supplying the gift sample of CGFZ as standard medicine. Additionally, the authors would like to acknowledge School of Pharmacy, ARKA JAIN University, Jamshedpur, Jharkhand worth granting access to the necessary research facilities.

References

- Sharma P., Kumar M., Sharma A., Arora D., Patial A., Rana M., 2019. An Overview on Green Chemistry. World J. Pharm. Pharm. Sci. 8(5), 202-208.



- Joshi D.R., Adhikari N., 2019. Green Chemistry: Beginning, Recent Progress, and Future Challenges. *World J. Pharm. Pharm. Sci.* 8(7), 280-293.
- Mane B.P., Antre V.R., Oswal J.R., 2012. Antidiabetic Drugs: An Overview. *Int. j. Pharm. Chem. Sci.* 1, 301-306.
- Balamurugan K., Mishra K., 2020. Optimization and Validation of the Simultaneous Determination of Vildagliptin and Metformin, in Bulk and Formulation by A Reverse Phase HPLC Method Using D-Optimal Experimental Design. *J. Glob. Pharma Technol.* 12(8), 1-12.
- Mishra K., Balamurugan K., Suresh R., 2018. Linagliptin: A Literature Review on Analytical and Bioanalytical Methods. *Int. J. Pharm. Qual. Assur.* 9(3), 225-230.
- Reddy M.R., Mishra K., Suresh R., 2018. Development and Validation of a Liquid Chromatographic Method for the Determination of Selected Anti Cancer Drugs in Bulk and Pharmaceutical Formulations. *Int. j. pharm. res. health sci.* 6(1), 2303-2307.
- Bhatt D., Thatavarthi P., Rajkamal B., 2018. Analytical Method Development and Validation for the Estimation of Canagliflozin in Bulk and Formulation by RP-HPLC. *Int. J. Pharm. Sci. Drug Res.* 10(3). 139-143.
- Sarangi B., Mishra K., Mohanta G.P., Manna P.K., 2019. In vitro-in vivo correlation (IVIVC) of solid lipid nanoparticles loaded with poorly water-soluble drug lovastatin. *Eur. Polym. J.* 122, 109366.
- Panigrahy U.P., Reddy A.S.K., 2015. A Novel Validated RP-HPLC-DAD Method for the Simultaneous Estimation of Metformin Hydrochloride and Canagliflozin in Bulk and Pharmaceutical Tablet Dosage Form with Forced Degradation Studies. *Orient. J. Chem.* 31(3), 1489-1507.
- Gurralla S., Shivaraj., Subrahmanyam C.V.S., Panikumar A.D., Gowthami S., 2019. Analytical Quality by Design Assisted HPLC Method for Quantification of Canagliflozin/ Metformin and Stability Studies. *Indian J. Pharm. Educ. Res.* 53(4), s699-s709.
- Mishra K., Dash A., Jabeen A., Vegesna S., Sahoo S.K., Gupta V., Jena D., 2023. Chemometric Assisted UV-Spectrophotometric Quantification of Tigecycline in Tablet Dosage Form. *Int. J. Drug Deliv. Technol.* 13(3), 976-981.
- Marella V.L., Syed A., Prasanna M.L., Nalluri B.N., 2017. A Novel Validated RP-HPLC Method for the Estimation of Canagliflozin in Bulk and Pharmaceutical Dosage Forms. *Int. J. Adv. Pharm. Anal.* 7(3), 24-27.
- Mishra K., Padmasri B., Vegesna S., Jabeen A., Dash A., Kumar S., Jena D., 2023. Chemometric Assisted UV-Spectrophotometric Quantification of Cefaclor in Suspension Dosage Form. *Int. J. Pharm. Qual. Assur.* 14(3), 734-739.
- Gautam M.K., Jena D., Jabeen A., Mukherjee R., Buralla K.K., Mishra K., 2023. Using Green Chemistry Concepts in Developing and Validating Analytical Methods for Meropenem in Parenteral Dosage Form: A Quality by Design Point of View. *J. Chem. Health Risks.* 13(6), 189-196.