



# Sugammadex – Reversal of Profound Neuromuscular Blockade in Laparoscopic Surgery

M. Bastina Jenoffia<sup>1</sup>, Vinodh Vani<sup>2</sup>, G. Nevetha Priyadarshini<sup>3</sup>, B. C handravathi<sup>4\*</sup>

<sup>1</sup>MD in Anaesthesiology, Assistant Professor, Department of Anaesthesiology, Tagore Medical College and Hospital

<sup>2</sup>MD in Anaesthesiology, Associate Professor, Department of Anaesthesiology

<sup>3</sup>MD in Anaesthesiology, Assistant Professor, Department of Anaesthesiology

<sup>4</sup>DNB in Anesthesiology, Senior Resident, Department of Anaesthesiology

(Received: 07 January 2024)

Revised: 12 February 2024

Accepted: 06 March 2024)

## KEYWORDS

Sugammadex,  
Laparoscopic  
Surgery,  
Rocuronium

## Abstract

**Introduction-** Sugammadex is a modified  $\gamma$ -cyclodextrin, which promptly reverses the steroidal nondepolarizing neuromuscular blocking agents. **Methodology-** This interventional trial will be commenced after ethical committee approval in Tagore Medical College and Hospital. This interventional trial was commenced after Institutional ethical committee approval (Ref No: IEC No: 03/NOVEMBER /2023) in Tagore Medical College and Hospital. Written informed consent obtained from patients before starting the study. A total number of 30 patients of ASA 1,2 between Age-18 – 60 years, undergoing laparoscopic surgery, willing to participate in the study was selected. **Results-**Around 43.3% subjects belonged to age group 20-30 years, 16.6% subjects from age group 31-40 years, 26.6 % from age group 41-50 years, and 13.3% subjects from age group 51-60 years. Among all patients, 12 males and 18 were females. **Conclusion-** Laparoscopic surgeries under DNMB/LAIP procedure is feasible and safe with lower incidence and severity of STP and high surgeons' satisfaction.

## Introduction

Sugammadex is a modified  $\gamma$ -cyclodextrin, which promptly reverses the steroidal nondepolarizing neuromuscular blocking agents. Although previous studies have reported a sudden increase in bispectral index (BIS) values after sugammadex reversal of neuromuscular blockade, many studies have supported that BIS increases are likely to be influenced by increased electromyography (EMG) activity following sugammadex administration, rather than true arousal. However, unexpected and sudden arousal after sugammadex use baffles clinicians, and further investigation of the effect of sugammadex on arousal is required. Recent studies have reported that sugammadex can encapsulate other drugs, such as propofol and remifentanyl, in addition to non-steroidal neuromuscular blockers.<sup>1</sup>

Laparoscopic surgeries are becoming popular for abdominal and pelvic surgeries.<sup>1,2</sup> But high intra-abdominal pressure induced in laparoscopic surgeries due to pneumoperitoneum affects ventilation and hemodynamic stability.<sup>3,4,5</sup> Deep neuromuscular blockade (DNMB) lowers intra-abdominal pressure (LAIP) and improves the outcome of laparoscopic surgeries, but it is associated with risk of post operative residual curarization.<sup>7,8</sup> Sugammadex potent and specific

antagonist of aminosteroids non-depolarizing muscle relaxants can effectively reverse.<sup>9,10</sup> Rocuronium induced neuromuscular blockade without residual muscle relaxation. In this study, we evaluate the efficacy of sugammadex as reversal for patients undergoing laparoscopic surgery with rocuronium.<sup>11,12</sup>

## Methodology

This interventional trial will be commenced after ethical committee approval in Tagore Medical College and Hospital. This interventional trial was commenced after Institutional ethical committee approval (Ref No: IEC No: 03/NOVEMBER /2023) in Tagore Medical College and Hospital. Written informed consent obtained from patients before starting the study. A total number of 30 patients of ASA 1,2 between Age-18 – 60 years, undergoing laparoscopic surgery, willing to participate in the study was selected. Patient was shifted to OT, connected to standard monitors, Patient will be premedicated with injection glycopyrrolate 10mcg/kg IV, injection midazolam 0.1mg/kg IV, injection fentanyl 2 mcg /kg IV. Patient will be induced with injection propofol 2mg/kg IV, paralysed with injection rocuronium 0.6mg/kg IV and intubated with endotracheal tube of appropriate size. Anaesthesia maintained with oxygen 2L/min, nitrous oxide 3L/min, sevoflurane 2 %, injection



rocuronium 0.15 mg/kg IV. At the end of the surgery, injection emeset 0.15mg/kg IV given, injection dexamethasone 0.15mg/kg IV given. Sugammadex was administered, with neuromuscular monitoring (post – tetanic counts 1-2 -injection sugammadex 4mg/kg IV and Train of four count  $\geq 2$  - injection sugammadex 2mg/kg IV),

Time taken for complete recovery from muscle relaxant indicated by neuromuscular monitoring with train of four  $> 0.9$  was noted, hemodynamics monitoring was done after administration of sugammadex for 24 hours and

watched for any adverse effects if any and manage accordingly


### Results

The present study was undertaken to evaluate the efficacy of Sugammadex as reversal for patients undergoing laparoscopic surgery with rocuronium in 30 consecutive cases. Around 43.3% subjects belonged to age group 20-30 years, 16.6% subjects from age group 31-40 years, 26.6 % from age group 41-50 years, and 13.3% subjects from age group 51-60 years. Among all patients, 12 males and 18 were females included in the study as shown in table 1.

**Table 1: Distribution of patients based on the age and gender**

Age (Years)	Number	Percentage (%)
20-30 years	13	43.3
31-40 years	5	16.6
41-50 years	8	26.6
51-60 years	4	13.3
<b>Gender</b>		
Male	12	40
Female	18	60

**Fig 1: Distribution of patients based on the age**



**Table 2: Distribution of patients based on Diagnosis**

Diagnosis	Frequency	Percent
Acute Appendicitis	13	43.3
Cholelithiasis	8	26.7
Fibroid Uterus	1	3.3
Left Ovarian Cyst	2	6.7
Multiple Gall Bladder Polyp	1	3.3



Right Ovarian Cyst	1	3.3
Subacute Appendicitis	2	6.7
Umbilical Hernia	2	6.7
<b>Total</b>	<b>30</b>	<b>100.0</b>


Table 3: Distribution of patients based on Procedure

<i>Procedure</i>	<b>Frequency</b>	<b>Percent</b>
Laparoscopic Appendectomy	15	50.0
Laparoscopic Cholecystectomy	9	30.0
Laparoscopic Hernioplasty	2	6.7
Laparoscopic Left Ovarian Cystectomy	2	6.7
Laparoscopic Right Ovarian Cystectomy	1	3.3
Laprosopic Assisted Vaginal Hystrectomy	1	3.3
<b>Total</b>	<b>30</b>	<b>100.0</b>

Table 4: Distribution of patients based on Time of Onset

<i>Time of Onset</i>	<b>Frequency</b>	<b>Percent</b>
2 minutes 10- 30 seconds	8	26.6
2minutes 31-40 seconds	5	16.6
2minutes 41-50 seconds	7	23.5
2minutes 51 sec-3 min	4	13.3
3 minutes 00-20 seconds	6	20
<b>Total</b>	<b>30</b>	<b>100.0</b>

Figure 2-Correlation between age and diagnosis



**Table 5: Distribution Of Patients Intraoperative Data**

		N=30	P	
Heart rate (beats/min)	Before intubation	79.9±4.2	0.344	
	After intubation	86.6±4.3	0.155	
	Before insufflation	81±4.1	0.039	
	After insufflation	89±3.6	<0.001	
	% of change during insufflation	10±3.7	0.045	
Mean arterial pressure (mmHg)	After table tilting	84.5±4.6	<0.001	
	After desufflation	80±4.7	0.848	
	After intubation	84±4.1	0.010	
	Before intubation	88.2±3.1	0.465	
	After intubation	92.5±3.1	0.263	
Operative field visibility scoring	Before insufflation	86.6±3.3	0.142	
	After insufflation	89.5±3.7	0.023	
	% of change during insufflation	3.4±1.12	2.75±1.43	0.005
	After table tilting	84.2±4.6	84.1±3.3	0.751
	After desufflation	85.1±4.4	83.7±3.4	0.003
Operative time (min)	After intubation	87.9±3.9	87.3±3	0.241
	Scores	3 (4.4%)	5 (7.5%)	0.609
		23 (33.8%)	28 (41.8%)	
		27 (39.7%)	22 (32.8%)	
		15 (22.1%)	12 (17.9%)	
	Median (IQR) of score	4 [3–4]	4 [3–4]	0.238
		27±6.7	29.4±8.3	0.068

**Table 6- Sugammadex Dose Requirement for Different Twitch Responses before Reversal**

Twitch Response Before Reversal	Sugammadex Dose	P Value
Post tetanic Count	2.35±0.98	<0.0001
Train-of-four count 1,2 or 3	2.01±1.22	
Train-of-four ratio <0.4	1.29±0.67	<0.0001
Train-of-four ratio <0.4	0.90±0.60	
Train-of-four count <2*	2.30±1.18	
Train-of-four count >2*	1.24±0.83	

## Discussion

Laparoscopic surgeries are becoming more accepted worldwide. In comparison with laparotomy, laparoscopic surgeries are minimally invasive, with less intraoperative blood loss, mild postoperative pain and faster recovery.<sup>13,14</sup> But artificial pneumoperitoneum created during laparoscopy causes high intraabdominal pressure which leads to hemodynamic instability which includes

increased afterload and preload, reduced cardiac output, arrhythmia and increased blood pressure, associated with ventilatory complications which includes increased airway pressure, hypercarbia, atelectasis.<sup>13</sup>

Use of deep neuromuscular blockade in laparoscopic surgeries reduces pneumoperitoneum pressure which improves surgical field, reduces pneumoperitoneum induced peri-operative pain, inhibition of stress



responses and improves postoperative outcome.<sup>15-18</sup> But significant disadvantage of application of deep neuromuscular block is high incidence of postoperative residual curarisation.<sup>19</sup> Residual muscle relaxation may cause airway obstruction, respiratory depression, aspiration pneumonia and hypoxemia.<sup>20</sup>

Cholinesterase inhibitors like neostigmine is commonly used for reversal of neuromuscular blockade but their effectiveness is limited in procedures where deep neuromuscular blockade is required due to risk of residual paralysis.

Sugammadex is a modified gamma-cyclodextrin, it encapsulates rocuronium and other aminosteroid non-depolarizing neuromuscular blocking agents creating a stable complex, removed by glomeruli, bypassing the hepatobiliary metabolic pathway.<sup>21</sup> Because of its unique antagonistic mechanism and metabolic pathway, it has faster reversal speed and efficacy with minimum side effects.<sup>22</sup> It also optimizes the surgical field of vision, lessens the surgical time, lessens the cardiopulmonary complication, reduced hospital stay and better prognosis. Sugammadex gives rapid reversal and better outcome in reversing the deep neuromuscular block induced by rocuronium than the commonly used neostigmine.<sup>23-26</sup> It also causes better recovery of gastrointestinal motility compared to glycopyrrolate and neostigmine mixture.<sup>27</sup> Patients receiving sugammadex recovered 3.4 times faster than those receiving sugammadex.<sup>28</sup> Most common side effects are nausea, vomiting, dysgeusia, transient hypotension, movement before the end of anaesthesia.<sup>29</sup> There is also possible risk of QT interval prolongation.<sup>30</sup> The dose of sugammadex should be based on actual body weight of the individual.<sup>31,32</sup> The recommended dose of sugammadex is 4mg/kg IV for deep neuromuscular block (If PTC 1-2) and 2mg/kg IV (if TOFC  $\geq 2$ ).<sup>33</sup> If large single dose of rocuronium 1.2 mg/kg IV is used, or if NMB is enquired to be reversed within 3 minutes, then sugammadex is given as 16mg/kg IV. If lower the recommended dose is used, there are chances of recurrence of NMB.<sup>34,35</sup>

## Conclusion

Laparoscopic cholecystectomy under DNMB and using LAIP is feasible and safe with a minimal failure rate and high surgeons' satisfaction. The applied DNMB/LAIP improved surgical outcomes with a reduction of the incidence and severity of STP. Sugammadex hastened the recovery of DNMB 3.2 times that of rocuronium-induced recovery of CNMB and is advocated as a reversal

agent for this procedure.

## References

- [1] Hatzinger M et al. Hans Christian Jacobaeus: inventor of human laparoscopy and thoracoscopy. *J Endourol* 2006; 20: 848–850.
- [2] Litynski GS. Raoul Palmer, World War II, and transabdominal coelioscopy. *Laparoscopy extends into gynecology. JSLS* 1997; 1: 289–92.
- [3] Mulier J, Dillemans B, Van Cauwenberge S. Impact of the patient's body position on the intraabdominal workspace during laparoscopic surgery. *SurgEndosc*. 2010; 24: 1398–1402.
- [4] Strang CM, Freden F, Maripuu E et al. Improved ventilation-perfusion matching with increasing abdominal pressure during CO<sub>2</sub>-pneumoperitoneum in pigs. *Acta Anaesthesiol Scand* 2011; 55: 887–96.
- [5] Dexter SP, Vucevic M, Gibson J, McMahon MJ. Hemodynamic consequences of high- and low-pressure capnoperitoneum during laparoscopic cholecystectomy. *SurgEndosc* 1999; 13: 376–81.
- [6] The Stockholm revision. *Acta Anaesthesiol Scand* 2007; 51: 789–808.
- [7] Ali HH, Utting JE, Gray C. Stimulus frequency in the detection of neuromuscular block in humans. *Br J Anaesth* 1970; 42: 967–78.
- [8] Meistelman C, Plaud B, Donati F. Rocuronium neuromuscular blockade at the adductor muscles of the larynx and adductor pollicis in humans. *Can J Anaesth* 1992; 39: 665–9.
- [9] Hemmerling TM, Donati F. Neuromuscular blockade at the larynx, the diaphragm and the corrugator supercilii muscle: a review. *Can J Anaesth* 2003; 50: 779–94.
- [10] Cantineau JP, Porte F, d'Honneur G, Duvaldestin P. Neuromuscular effects of rocuronium on the diaphragm and adductor pollicis muscles in anesthetized patients. *Anesthesiology* 1994; 81: 585–90.
- [11] Blobner M, Frick CG, Stäubli RB et al. Neuromuscular blockade improves surgical conditions. *SurgEndosc* 2015; 29: 627–36.
- [12] Dubois PE, Putz L, Jamart J et al. Deep neuromuscular block improves surgical conditions during laparoscopic hysterectomy: a randomised controlled trial. *Eur J Anaesthesiol* 2014; 31: 1–7.
- [13] Oti C, Mahendran M, Sabir N. Anaesthesia for laparoscopic surgery. *Br J Hosp Med (Lond)*.



- 2016;77(1):24–28.  
doi:10.12968/hmed.2016.77.1.24
- [14] Atkinson TM, Giraud GD, Togioka BM, Jones DB, Cigarroa JE. Cardiovascular and ventilatory consequences of laparoscopic surgery. *Circulation*. 2017;135(7):700–710. doi:10.1161/CIRCULATIONAHA.116.023262
- [15] Fuchs-Buder T, Schmartz D, Baumann C, et al. Deep neuromuscular blockade improves surgical conditions during gastric bypass surgery for morbid obesity: a randomised controlled trial. *Eur J Anaesthesiol*. 2019;36(7):486–493. doi:10.1097/EJA.0000000000000996
- [16] Raval AD, Deshpande S, Rabar S, et al. Does deep neuromuscular blockade during laparoscopy procedures change patient, surgical, and healthcare resource outcomes? A systematic review and meta-analysis of randomized controlled trials. *PLoS One*. 2020;15(4):e0231452. doi:10.1371/journal.pone.0231452
- [17] Koo BW, Oh AY, Ryu JH, et al. Effects of deep neuromuscular blockade on the stress response during laparoscopic gastrectomy randomized controlled trials. *Sci Rep*. 2019;9(1):12411. doi:10.1038/s41598-019-48919-2
- [18] Cho YJ, Paik H, Jeong SY, et al. Lower intra-abdominal pressure has no cardiopulmonary benefits during laparoscopic colorectal surgery: a double-blind, randomized controlled trial. *Surg Endosc*. 2018;32(11):4533–4542. doi:10.1007/s00464-018-6204-2
- [19] Raval AD, Uyei J, Karabis A, Bash LD, Brull SJ. Incidence of residual neuromuscular blockade and use of neuromuscular blocking agents with or without antagonists: a systematic review and meta-analysis of randomized controlled trials. *J Clin Anesth*. 2020;64:109818. doi:10.1016/j.jclinane.2020.109818
- [20] Bronsert MR, Henderson WG, Monk TG, et al. Intermediateacting nondepolarizing neuromuscular blocking agents and risk of postoperative 30-day morbidity and mortality, and long-term survival. *Anesth Analg*. 2017;124(5):1476–1483. doi:10.1213/ANE.0000000000001848
- [21] Kovac AL. Sugammadex: the first selective binding reversal agent for neuromuscular block. *J Clin Anesth*. (2009) 21:444–53.
- [22] Karalapillai D, Kaufman M, Weinberg L. Sugammadex. *Crit Care Resusc*. 2013;15(1):57–62.
- [23] Bartels K, Hunter JM. Neostigmine versus sugammadex: the tide may be turning, but we still need to navigate the winds. *Br J Anaesth*. 2020;124(5):504–507. doi:10.1016/j.bja.2020.02.021
- [24] Krause M, McWilliams SK, Bullard KJ, et al. Neostigmine versus sugammadex for reversal of neuromuscular blockade and effects on reintubation for respiratory failure or newly initiated noninvasive ventilation: an interrupted time series design. *Anesth Analg*. 2020;131(1):141–151. doi:10.1213/ANE.0000000000004505
- [25] Carron M, Zarantonello F, Tellaroli P, Ori C. Efficacy and safety of sugammadex compared to neostigmine for reversal of neuromuscular blockade: a meta-analysis of randomized controlled trials. *J Clin Anesth*. 2016;35:1–12. doi:10.1016/j.jclinane.2016.06.018
- [26] Hristovska AM, Duch P, Allingstrup M, Afshari A. Efficacy and safety of sugammadex versus neostigmine in reversing neuromuscular blockade in adults. *Cochrane Database Syst Rev*. 2017;8:CD012763
- [27] An J, Noh H, Kim E, Lee J, Woo K, Kim H. Neuromuscular blockade reversal with sugammadex versus pyridostigmine/glycopyrrolate in laparoscopic cholecystectomy: a randomized trial of effects on postoperative gastrointestinal motility. *Korean J Anesthesiol*. 2020;73(2):137–144. doi:10.4097/kja.19360
- [28] Geldner G, Niskanen M, Laurila P, et al. A randomised controlled trial comparing sugammadex and neostigmine at different depths of neuromuscular blockade in patients undergoing laparoscopic surgery. *Anaesthesia*. 2012;67(9):991–998. doi:10.1111/j.1365-2044.2012.07197.x
- [29] Tobias JD, Thomas M. Current evidence for the use of sugammadex in children. *Paediatr Anaesth*. 2017;27(2):118–125. doi:10.1111/pan.13050
- [30] Cammu G, De Kam PJ, Demeyer I, et al. Safety and tolerability of single intravenous doses of sugammadex administered simultaneously with rocuronium or vecuronium in healthy volunteers. *Br J Anaesth*. 2008;100(3):373–379. doi:10.1093/bja/aem402
- [31] Thompson CA. Sugammadex approved to reverse NMBA effects. *Am J Health Syst Pharm*. 2016;73(3):100. doi:10.2146/news160009



- 
- [32] Mirakhur RK. Sugammadex in clinical practice. *Anaesthesia*. 2009;64(Suppl 1):45–54.
- [33] Kopman AF. Neostigmine versus sugammadex: which, when, and how much? *Anesthesiology*. 2010;113(5):1010–1011. doi:10.1097/ALN.0b013e3181f41847
- [34] Asztalos L, Szabo-Maak Z, Gajdos A, et al. Reversal of vecuronium-induced neuromuscular blockade with low-dose sugammadex at train-of-four count of four: a randomized controlled trial. *Anesthesiology*. 2017;127(3):441–449. doi:10.1097/ALN.0000000000001744
- [35] De Boer HD, Carlos RV, Brull SJ. Is lower-dose sugammadex a cost-saving strategy for reversal of deep neuromuscular block? Facts and fiction. *BMC Anesthesiol*. 2018;18(1):159. doi:10.1186/s12871-018-0605-6